

On the Origin of Magnetic Fields in GRB Sources

Eric G. Blackman

*Dept. of Physics & Astronomy & Lab. for Laser Energetics, Univ. of Rochester
Rochester, NY, 14627 USA*

Abstract

Magnetic fields play a dual role in gamma-ray bursts (GRBs). First, GRB and afterglow spectra (the latter interpreted as emission from external shocks) imply synchrotron radiation in a magnetic field that is a significant fraction of equipartition with the particle energy density. Second, magnetized rotators with $\sim 10^{15}$ Gauss field may power GRB by transporting Poynting flux to large distances where it dissipates and also drives an external shock. The field amplification at external shocks and in the engine involve separate processes. External shock fields are likely either seeded by a pre-GRB wind, or are amplified by two-stream plasma instabilities with MHD turbulence playing a subsequent role. In the engine, the large scale fields are likely produced by MHD helical dynamos, since flux accretion cannot easily compete with turbulent diffusion, and because structures must be large enough to rise to coronae before diffusing. Why helical dynamos are feasible, and their relation to the magnetorotational instability are among the points discussed.

1 Introduction

Magnetic fields are an intermediary between gravity and radiation. Often their GRB consequences are explored without addressing their origin. Here I focus on the latter.

Consider GRB shocks [1, 2]: external shocks, driven by relativistic outflows acting as a piston, propagate into the external medium and accelerate particles that contribute to afterglow emission. In contrast, internal shocks would occur inside the ejecta and have been invoked to explain the GRB itself. Field amplification at the external shock poses the same problem whether or not the ejecta is Poynting flux or gas driven, but internal shocks apply only to hydrodynamic models.

Despite their advantages, Poynting flux models [3, 4, 5] raise questions of how the emission is produced [6], and where the strong ordered field comes from. In situ helical dynamo amplification [4, 7, 8] is well suited for amplifying the large scale fields required for producing the inferred jets of GRB [9], particularly in light of recent understanding of nonlinear dynamos [8, 10, 11, 12]. Nonlinear helical dynamos can operate in a convectively driven star or in a stratified disk [15, 16] unstable to the magnetorotational instability (MRI) [17, 18].

2 Field amplification at external shocks

Magnetic fields are likely generated as part of relativistic collisionless GRB shock formation. Both observations and modeling suggest a magnetic to particle energy density ratio $10^{-5} < \epsilon_m/\epsilon_{part} \simeq 10^{-1}$ (c.f. [2]). The ambient field of a typical surrounding ISM is $\sim 10^{-6}$ Gauss. Compression of the ambient field from a relativistic shock (of Lorentz factor γ) would give $\sim 10^{-4}(\gamma/10^2)(B_{ext}/10^{-6}\text{G})\text{G}$, but the equipartition value is $B_{eq} \sim 10^2(\gamma/10^2)(n/1\text{cm}^{-3})^{1/2}\text{G}$, so an amplification mechanism is needed.

The two-stream Weibel instability in the GRB context [19] may be important. Here the field draws energy from free-streaming charged particles. Imagine a weak seed magnetic field whose amplitude has a sinusoidal perturbation in the plane perpendicular to the flow direction. The resulting magnetic force guides charged particles of one sign toward each other on one side of the perturbation plane and away from each other on the opposite side of the perturbation. This creates current channels which amplify the field by induction, further amplifying the currents, and ultimately exponentially amplifying the field. Initially anisotropic particle distributions become isotropic. The magnetic fields gets the randomized energy, which also helps the shock form. The instability saturates first on electrons and then on protons, such that the proton gyro-radius approaches the relativistic Debye length (or skin depth) $\gamma_{th}^{1/2}c/\omega_p$ (the scale of charge separation, where γ_{th} is the thermalized Lorentz factor and ω_p is the plasma frequency). Saturation could produce energy equipartition between protons and the magnetic field.

The above saturation scale would be ~ 8 orders of magnitude smaller than the source emission region. How subsequent field growth/advection evolves needs more work. A few simulations [20] show that regions of field amplification do seem to convect beyond the skin depth, but longer simulations are needed. It may actually take some distance before the full flow cross section participates in the instability, after which the field evolution is unclear. Perhaps MHD instabilities (super-gyroradius scales) can grow and further amplify the field over larger scales as in supernovae [21].

Another alternative: some engine models [22, 23] involve multiple winds in the context of massive stellar collapse. The core first collapses first to a neutron star (NS), and then later to a black hole (BH). The GRB external shock could then propagate into a pulsar wind “pre-magnetized” bubble [23]. Outside of the light cone, the field falls off as $B \sim B_s \left(\frac{R_s^3}{r_{lc}^2 r}\right) \sim 10^2 \left(\frac{R}{10^6\text{cm}}\right) \left(\frac{r}{10^{15}\text{cm}}\right)^{-1} \left(\frac{B_s}{10^{16}\text{G}}\right) \left(\frac{\Omega}{10^4\text{s}^{-1}}\right)^2\text{G}$, where B_s is the surface field, r_{lc} is the light cone, R_s is the scale of the rotator and r is the radius.

3 Cold, poynting flux GRB models

Small scale magnetic fields dissipating in the engine could in principle drive a hot jet. But collimation in such models is hard to understand, and copious radiation would

be produced. The outflow then has to convert this energy to bulk motion and then back to radiating particles. Perhaps more appealing is the cold, baryon free, Poynting flux paradigm [3, 4, 5, 14] in which particle acceleration occurs only at large distances where flux freezing is broken (and where the acceleration is needed). The magnetic hoop stress can provide jet collimation.

Poynting flux outflows are actually powered by rotation; the magnetic field acts as a drive belt and determining the luminosity. A large fraction of the rotational energy of the anchoring compact object can in principle be used (though we don't fully understand the dissipation mechanism yet [6].) This may differ from internal shock models, where only the relative energy between colliding blobs in the flow is used.

Poynting flux GRB rotators can arise in non-supernova events [(i) magnetar formed from accretion induced collapse of a white dwarf [3, 4]. (ii) merger that produces a NS/torus + BH [5]] or in supernova events [(i) NS $\rightarrow$ disk + BH from fall-back of material after breakout in a partially failed or weak SN, with GRB powered by accretion [24], (ii) or powered by NS rotation [3, 6, 13, 14] (iii) or delayed BH jet formation after supermassive NS forms in a supernova (=supranova) [22] with a BH accretion driven magnetic outflow propagating into an earlier pulsar wind [23] (which could pre-magnetize the ambient medium for external shocks.)] In each of these cases exponential field amplification (rather than linear [13]) can occur, either in the NS torus, in the NS itself [4] or at NS boundary by MRI [25] or helical interface dynamo as in an AGB star [26] But these dynamo proposals must be evaluated in light of recent work on nonlinear helical and nonhelical dynamos (section 5).

The basic magnetic outflow launching is either a “spring” [27] driven by toroidal field pressure in a corona of a disk or outside the light cylinder of a magnetar, or a centrifugal “fling” [28] of material along poloidal field which develops a subsequent toroidal magnetic pressure. The maximum luminosity in either case is $\sim B^2 r^3 \Omega$ with quantities measured at the inner most distance r at which the toroidal field becomes of order the poloidal field: e.g. near the surface of an accretion disk, or at the light cylinder for a magnetar. In the latter, the luminosity formula becomes the dipole radiation formula when scaled to the stellar surface.

4 Need for in situ generation of large scale fields

4.1 Small scale fields shred before they escape

For magnetic fields to launch jets or power coronae, the field buoyancy time must be shorter than the diffusion time in the disk. The buoyancy time is $\sim h/\overline{U}_A$ where h is the system thickness scale and $\overline{U}_A$ is the Alfvén speed associated with the structure of scale L . To escape without diffusing, $h/\overline{U}_A < L^2/\beta$ where the turbulent magnetic diffusion coefficient $\beta \sim (u_2 l)$ with u_2 and l the dominant turbulent speed and scale. (e.g. $l \sim$ cube root of the turbulent cell volume $(l_x l_y l_z)^{1/3}$). Escape then

requires $L^2 > lh(u_2/\bar{U}_A)$. The ratio on the right is typically $\gtrsim 1$ (more on that later) so $L^2 > lh$ is required for escape.

Now consider a Shakura-Sunayev [29] disk viscosity $\nu_{ss} \sim \beta = u_2 l \sim \alpha_{ss} c_s h \sim u_2^2/\Omega \sim u_{A,2}^2/\Omega$, where $u_{A,2}$ is the Alfvén speed associated with the turbulent B -field and $u_2 \sim u_{A,2}$ follows from simulations [18, 15]. Then $\alpha_{ss} \sim u_{A,2}^2/c_s^2$ or $u_{A,T} \sim \alpha_{ss}^{1/2} c_s$, so that $l = \alpha_{ss}^{1/2} h$. The escape condition $L^2 > lh$ then gives $L > \alpha_{ss}^{1/4} h$ *smallest* dimension of the *largest* structures since the former determines the diffusion time. The escape condition is not satisfied in the nonhelical MRI dynamo (see maximum k_y, k_z in fig. 4 of [18]). Note also if $L = l \sim \alpha^{1/2} H$, then escape requires $\alpha_{ss} \sim 1$.

4.2 Accretion of flux can’t easily beat turbulent diffusion

Can the large scale fields required by jets be accreted onto a central engine? Though 3-D MHD simulations are needed, the answer seems to be “no” particularly for thin disks. The reason is that MHD the turbulence produces a turbulent magnetic diffusivity $\beta \sim \nu_{ss}$. To determine if a vertical field can be accreted in an MHD turbulent disk, the ratio of rates of flux advection to diffusion is needed. The dominant variation of the field strength is likely in the vertical direction across the thickness of the disk and the dominant variation of the radial velocity is in the r direction. The above mentioned ratio is then $\nabla \times (\bar{\mathbf{v}} \times \bar{\mathbf{B}})/(\beta \nabla^2 \bar{\mathbf{B}}) \sim (h^2/R^2)(\alpha_{ss} h c_s/\beta) \sim h^2/R^2$. The diffusion term thus dominates for a thin disk. This was shown with a numerically 2-D calculation in [30]. and motivates an in situ helical dynamo which can overcome diffusion.

5 MHD in situ amplification

It is useful to categorize dynamos into two classes: direct and inverse. Direct dynamos sustain magnetic fields on scale at or below the dominant turbulent scale. They do not require helical turbulence although helicity does influence the resulting magnetic spectrum. In contrast, an inverse dynamo describes amplification on scales larger than that of the dominant scale of the turbulence. The label “inverse” is used to suggest an inverse cascade. The turbulence must be helical to generate and sustain large scale flux over times longer than an eddy turnover time in order to drive and collimate jets.

5.1 Direct dynamo

In a nonhelical direct dynamo, the field grows by line stretching in a turbulent random walk. The turbulence can be externally driven by isotropic forcing, or self-generated by an angular velocity gradient. In either case, turbulent stretching compensates for

exponential decay from turbulent diffusion. The latter operates because the random motions of the gas also mix the field lines, inducing a cascade to small scales where dissipation occurs. In 3-D, a steady state balance can be achieved. In the saturated state, the magnetic energy $\sim$ turbulent kinetic energy below the dominant turbulent scale, but the spectral shape is a subtle issue.

5.1.1 Forced turbulence

In numerical experiments where turbulence is forced isotropically in a periodic box without helical correlations, the field piles up at the resistive scale [31, 32] for isotropic viscosity and for magnetic Prandtl number $\equiv \nu/\lambda > 1$ (where ν is the microphysical viscosity and λ is the magnetic diffusivity). When the turbulence is forced with a fractional kinetic helicity $f_h \sim \langle \mathbf{u} \cdot \nabla \times \mathbf{u} \rangle / k_2 u^2 > 0.4$ (k_2 is the forcing wavenumber ~ 5 in Refs. [10, 33]), the direct dynamo then piles up magnetic energy at the forcing scale of the turbulence, not the resistive scale [33]. (A second peak also develops at larger scales due to the inverse dynamo discussed in section 5.)

For $f_h = 0$, $Pr > 1$, and isotropic viscosity, a nonlocal effect induces the field pile up on small scales [32, 34]: forcing scale motions directly stretch the field into long magnetic folds which can't unwind for $Pr_M > 1$

5.1.2 Nonstratified MRI: direct nonhelical dynamo, no sustained flux

In a disk (or star) with a weak magnetic field, the MRI is driven by the angular velocity gradient and is thus different from isotropically forced turbulence. In the absence of stratification, and thus the absence of a pseudoscalar helicity, the MRI does not produce magnetic fields on scales larger than the scale of the developed turbulence. The largest scale of the field in the toroidal direction does approach the scale of the box height in simulations [18], but the kinetic turbulence also approaches this scale. The marginally unstable (and dominant) growth mode of the MRI satisfies $ku_A \sim \Omega$ [17] so that as the magnetic field is amplified from the instability, larger and larger scales become unstable to growth. The largest scale that can grow is that associated the scale height h . Without stratification or helicity, one would expect the largest scale of the turbulence to be of order the largest scale of the field and this is what is seen (Fig. 4 of [18] where k_y is the toroidal direction.)

Note that the MRI, unlike the isotropic forced turbulence, produces a nearly Kolmogorov spectrum in the field even without any helicity. The role of large scale shear (rather than helicity) in keeping the field from piling up on the resistive scale needs to be further investigated. That being said, note that the flux associated with the field on the largest scale in the nonstratified MRI would change every turnover time as the field is shredded. The field is large scale in the nonstratified MRI only in the sense that the box height limits the scale of the turbulence NOT because there is

any sustained ordered flux on scales larger than the turbulence. Furthermore, when the box is periodic, the net flux cannot grow, as it is conserved. Note also that the scale of the largest field structure in the vertical and the radial (k_x) directions are smaller than in the toroidal direction. This is important in the context of section 4.1. Because of the rapid change in sign of the flux and the constraints of section 4.1, a nonstratified nonhelical MRI case would *not* produce the fields required for jets— but real rotators do have stratification and a helical version would work (see sect. 5.2.3).

5.2 Helical inverse dynamo

The helical inverse dynamo amplifies field on scales larger than that of the turbulence. This dynamo is the most plausible for the production of large scale fields for cold Poynting flux GRB models.

Here is the basic mechanism [7, 35]: Consider an initially weak toroidal (=encircling the rotation axis) loop of the magnetic field embedded in the rotator. With an outwardly decreasing density gradient a rising turbulent eddy threaded by a magnetic field will twist oppositely to the underlying global rotation to conserve angular momentum. Statistically, northern (southern) eddies twist the field clockwise (counterclockwise). This is the “ α_d ” effect and the result is a large scale poloidal field loop. Differential rotation shears this loop (the “ Ω ”-effect). The bottom part reinforces the initial toroidal field and the top part diffuses away. The result is exponential growth. Inside a star the α_d effect can be supplied by neutrino driven convection + a vertical density gradient. For an MRI driven system, such stratification is also required.

This is revealed mathematically by averaging the magnetic induction equation over a local volume and breaking all quantities (velocity $\mathbf{U}$, magnetic field $\mathbf{B}$ in Alfvén velocity units, and normalized current density $\mathbf{J} \equiv \nabla \times \mathbf{B}$) into their mean (indicated by an overbar) and fluctuating (lower case) components. The result is [7]:

$$\partial_t \overline{\mathbf{B}} = \nabla \times (\alpha_d \overline{\mathbf{B}} + \overline{\mathbf{U}} \times \overline{\mathbf{B}}) + (\beta + \lambda) \nabla^2 \overline{\mathbf{B}}. \quad (1)$$

The $\overline{\mathbf{U}}$ term incorporates the Ω -effect, the β term incorporates the turbulent diffusion (assume constant β) and the first term on the right incorporates the α_d -effect. The quantity α_d is given by [12, 36] $\alpha_d = \alpha_{d0} + (\tau/3) \overline{\mathbf{b} \cdot \nabla \times \mathbf{b}}$, where $\alpha_{d0} = -(\tau/3) \overline{\mathbf{u} \cdot \nabla \times \mathbf{u}}$, τ is a turbulent damping time and $\overline{\mathbf{u} \cdot \nabla \times \mathbf{u}}$ is the kinetic helicity. In textbooks, $\alpha_d = \alpha_{d0}$, but properly including the magnetic forces that backreact on the velocity driving the field growth lead to the above correction term. This is studied formally in Ref. [12].

From the form of α_d , it is evident that the correction term can offset α_{d0} and quench the dynamo [11, 37, 12]. The principle of magnetic helicity conservation determines the extent to which α_d is suppressed. The magnetic helicity, a volume integral $H \equiv \int \mathbf{A} \cdot \mathbf{B} dV \equiv \langle \mathbf{A} \cdot \mathbf{B} \rangle V$ satisfies [38, 39]

$$\partial_t H = -2\lambda C - \text{surface terms}, \quad (2)$$

where $\mathbf{A}$ is defined by $\mathbf{B} = \nabla \times \mathbf{A}$, and the current helicity C is defined by $C \equiv \langle \mathbf{J} \cdot \mathbf{B} \rangle V$. Without non-diffusive surface terms H is well conserved: for $\lambda \rightarrow 0$ the λ term in (2) converges to zero [40]. Since H is a measure of “linkage” and “twist” of field lines [39], its conservation implies that the α_d -effect does not produce a net magnetic twist, just positive and negative magnetic twists on different scales. [11, 41].

The importance of the scale segregation is most easily seen in a two-scale approach. Write $H = H_1 + H_2$, where $H_1 = \langle \overline{\mathbf{A}} \cdot \overline{\mathbf{B}} \rangle V$ and $H_2 = \langle \overline{\mathbf{a}} \cdot \overline{\mathbf{b}} \rangle V$ correspond to volume integrated large and small scale contributions respectively. Also, $C = \langle \overline{\mathbf{J}} \cdot \overline{\mathbf{B}} \rangle V + \langle \overline{\mathbf{j}} \cdot \overline{\mathbf{b}} \rangle V = k_1^2 H_1 + k_2^2 H_2$, where k_1 and k_2 represent the wavenumbers (inverse gradients) associated with the large and small scales respectively, and the second equality follows for a closed system.

Now relate $\overline{\mathbf{B}}$ to H_1 . Define ϵ_1 such that the large scale magnetic energy $\langle \overline{\mathbf{B}}^2 \rangle V = H_1 k_1 / \epsilon_1$ and where $0 < |\epsilon_1| \leq 1$, where $|\epsilon_1| = 1$ only for a force-free helical large scale field (i.e. for which $\overline{\mathbf{J}} \parallel \overline{\mathbf{B}}$ so that the force $\overline{\mathbf{J}} \times \overline{\mathbf{B}} = 0$). In a northern hemisphere $\epsilon_1 > 0$. The conservation equations analogous to (2) for H_1 and H_2 are

$$\partial_t H_1 = 2(\langle \alpha \rangle k_1 / \epsilon_1 - \langle \beta \rangle k_1^2) H_1 - 2\lambda k_1^2 H_1 - \text{surface terms} \quad (3)$$

and

$$\partial_t H_2 = -2(\langle \alpha \rangle k_1 / \epsilon_1 - \langle \beta \rangle k_1^2) H_1 - 2\lambda k_2^2 H_2 - \text{surface terms} \quad (4)$$

where $\langle \alpha \rangle = (\langle \alpha_0 \rangle + \frac{1}{3} \tau k_2^2 H_2 / V)$. The case without non-diffusive surface terms and with $\epsilon_1 = 1$ is called an α_d^2 dynamo. The solution [11, 12, 37]. shows that for initially small H_2 but large α_0 , H_1 grows. Growth of H_1 implies the oppositely signed growth of H_2 . This H_2 backreacts on α_0 , ultimately quenching $\langle \alpha_d \rangle$ and the dynamo. The theory quantitatively matches simulations of [10]. At early times, kinematic growth is unimpeded, and the large scale field can grow to $\overline{U}_A^2 \sim f_h(k_1/k_2) u_2^2$, where f_h is the fractional kinetic helicity. (In general $f_h \leq 1$ must be estimated, but $f_h \sim 1$ will be later derived for young NS/torus engines of GRB.) Eventually, the small-scale magnetic helicity backreacts on the kinetic helicity, suppressing the growth rate to a resistively limited value. Ultimate saturation occurs at $\overline{U}_A^2 \simeq f_h(k_2/k_1) u_2^2$ but growth beyond the kinematic regime is too slow to be of relevance for GRBs.

5.2.1 Role of the Rossby Number R_o

When differential rotation operates over the same scale as α_d , a rotation to eddy turnover time ratio $R_o \equiv t_{rot}/t_{ed} = v_2 k_2 / \Omega > 1$ favors an α_d^2 dynamo over an $\alpha_d - \Omega$ dynamo. This follows from comparing the Ω and α_d terms effects in the $\partial_t \overline{\mathbf{B}}_\phi$ equation. But the Ω layer could be $\ll$ than the α_d layer [26]. Then R_o is not the determining parameter, but either dynamo produces the desired large scale fields. A more important role of R_o : for the growth term in (3) to overcome decay, $|\epsilon_1| \beta k_1 / \alpha_{d0} \sim |\epsilon_1| R_o < 1$, using $k_1 h \sim 1$, $\beta \sim u_2 / k_2$ and using $\alpha_{d0} \sim \Omega \cdot \nabla \rho / \rho k_2^2 \sim \Omega / h k_2^2$ [7, 35] for the case $R_o \gtrsim 1$. Since $|\epsilon_1| \leq 1$ the condition can be met.

5.2.2 Application to young NS/torus GRB progenitors

Here I cavalierly apply the nonlinear results of sect. 5.2 to young NS-like MHD rotators typical of Poynting flux GRB engine models [4]. Without buoyancy, the amplification term in the kinematic regime is $\sim \alpha_{d0}k_1 - \beta k_1^2$. Although buoyancy dominates the β term when $1/k_1\bar{U}_A < 1/\beta k_1^2$ (where k_1 is used for the inverse system size and the wavenumber of $\bar{B}$) for $f_h \sim 1$, it does not limit $\bar{U}_A$ calculated in sect. 5.2 because a buoyancy time $H/\bar{U}_A$ equals a growth time when $f_h v_2 \sim \bar{U}_A$, which is $(k_2/k_1)^{1/2}$ larger than the saturation value from mag. helicity conservation for $f_h = 1$.

The f_h for the young NS case can be estimated from $f_h \sim \alpha_{d0}/u_2$ taking $u_2 \sim \times 10^8 F_{39} \text{cm/s}$ as the neutrino driven turbulent convection velocity, and F_{39} is the neutrino heat flux in units $10^{39} \text{ erg/sec cm}^{-2}$ [4]. Using α_{d0} from Ref. [7] and the NS numbers [4], $\alpha_d \sim 10^8 \left(\frac{\Omega}{6 \times 10^3 \text{sec}^{-1}} \right) \left(\frac{l}{2 \times 10^5 \text{cm}} \right)^2 \left(\frac{h}{10^6 \text{cm}} \right)^{-1} \text{cm/s}$, so $f_h \sim 1$. From section 5.2 the maximum field of a closed system at the end of the kinematic regime is then $\bar{U}_A = u_2(k_1/k_2)^{1/2}$. As applied to the young NS convection zone, u_2 corresponds to the convective equipartition field B_{eq} for $f_h \sim 1$, and for $k_1/k_2 \sim 1/5$ we then have $\bar{B} \sim (k_1/k_2)^{1/2} B_{eq} \sim 4 \times 10^{15} F_{39} \text{G}$, with growth time is $\sim 10^{-3} \text{sec}$. This is sufficient for GRB. This estimate from an α_d^2 dynamo is a lower limit to that from an $\alpha_d - \Omega$ dynamo since the latter gains from shear, such that $|\epsilon_1| < 1$ [37]. But since $f_h \sim 1$ corresponds to $|\epsilon_1|R_o \sim k_1/k_2$, for $R_o \sim 1$, the lower limit $\sim$ actual value.

5.2.3 Helical MRI dynamo

In a sheared rotator, the presence or absence of an MRI dynamo *does not* depend on helicity but the additional presence of an inverse helical dynamo (described in section 5.2) does. Only a helical dynamo can sustain a large scale ordered magnetic field and magnetic flux over long time scales. This has been demonstrated in stratified shearing box simulations [16]. There MRI turbulence can generate its own kinetic or current helicity producing a finite α_d . A global large scale flux is also produced if periodic boundary conditions are avoided. Figure 2 of [16] clearly shows the effect of a helical dynamo operating in an MRI simulation, where a large scale field with cycle period of 30 orbit periods is seen. Thus the large scale fields that satisfy the escape condition of section 4.1 can in general also be produced by the MRI, but only when density stratification is included. Using the MRI in spherical geometry [42] to generate field at the boundary of the NS GRB progenitor [25] should involve the MRI as a potential source (in addition to convection) of a helical dynamo, not just a nonhelical one.

6 Summary points

(1) The problem of field amplification in external shocks is the same for both hot and Poynting flux engine models. Weibel instability + MHD turbulence may work in

succession. (2) Large scale fields in the engine are required for GRB cold Poynting flux jets and are difficult to accrete because of turbulent diffusion. At minimum, flux accretion requires very thick $h \sim R$ disks. (3) Buoyancy must beat turbulent diffusion to get field in coronae to produce jets. Even this requires large scale fields. (4) A balance between helical dynamos and buoyancy can produce the required GRB engine fields. The expected large scale field strength is proportional to the ratio of turbulent to system scale for fully nonlinear helical dynamos. Strong large scale fields result. (6) The magnetorotational instability (MRI) can lead to a nonhelical or helical dynamo, but only the helical version (expected in nature's stratified, sheared rotators) favorably produces the large scale fields required for jets.

Thanks to B. Paczynski and V. Pariev for recent discussions, and to the Dept. of Astrophysical Sciences, Princeton Univ. for hospitality.

References

- [1] M.J. Rees & P. Mészáros, MNRAS **258**, 41P (1992); M.J. Rees, & P. Meszaros, ApJL, **430**, L93 (1994); R. Sari & T. Piran, MNRAS, **287**, 110 (1997); E. Waxman, ApJL **489**, L33 (1997)
- [2] T. Piran, Phys. Rep, **333** 529 (2000) ; P. Mészáros, ARAA, **40**, 137 (2002)
- [3] V.V. Usov, Nat., **357**, 472 (1992); E.G. Blackman, I. Yi, & G.B. Field, ApJL **473**,L79 (1996)
- [4] R.C. Duncan & C. Thompson, ApJL, **392**, L9 (1992); C. Thompson & R.C. Duncan, ApJ, **408**, 194 (1993);
- [5] P. Meszaros, & M.J. Rees, ApJL, **482**, L29 (1997)
- [6] E.G. Blackman & I. Yi, ApJL, **498**,L31 (1998); M. Lyutikov,& E.G. Blackman, MNRAS **321**,177 (2001)
- [7] H.K. Moffatt, 1978, Magnetic Field Generation in Electrically Conducting Fluids (Cambridge University Press, Cambridge)) E.N. Parker 1979, Cosmical Magnetic Fields (Clarendon Press, Oxford);
- [8] E.G. Blackman (astro-ph/0205002) and A. Brandenburg, (astro-ph/0207394) articles to appear in Turbulence and Magnetic Fields in Astrophysics eds. E. Falgarone and T. Passot, Springer Lecture Notes in Physics (2002),
- [9] J.E. Rhoads, ApJ, **487**,L1 (1997) R. Sari, T. Piran, & J.P. Halpern, ApJL, **519**,L17 (1999); J. Granot, A. Panaitescu, P. Kumar, & S.E. Woosley, ApJL, **570**, L61 (2002);E. Berger et al., ApJ, **556**, 556 (2001); G. Björnsson et al., J. Hjorth, K. Pedersen, & J.U. Fynbo, ApJL **579**, L59 (2002);

- [10] A. Brandenburg, ApJ, **550**,824 (2001)
- [11] G.B. Field & E.G. Blackman, ApJ, **572**, 685 (2002)
- [12] E.G. Blackman and G.B. Field, PRL, in press (2002).
- [13] M.A. Ruderman, L. Tao, L., & W. Kluźniak ApJ **542**, 243 (2000)
- [14] J.C. Wheeler, I. Yi, P. Höflich, P., & L. Wang, ApJ, **537** 810 (2000)
- [15] A. Brandenburg, A. Nordlund,, R.F. Stein,& U. Torkelsson, ApJ **446** 741 (1995)
- [16] A. Brandenburg & K.J. Donner, MNRAS **288** L29 (1997)
- [17] S.A. Balbus & J.F. Hawley, ApJ, **376**, 214 (1991); S.A. Balbus & J.F. Hawley
Rev Mod Phys., **70**, 1 (1998)
- [18] J.F. Hawley, C.F. Gammie, & S.A. Balbus, ApJ **440**, 742 (1995)
- [19] M.V. Medvedev & A. Loeb, ApJ **526** 697 (1999)
- [20] A. Nordlund, this volume; F. Califano, F., T. Cecchi, & C. Chiuderi, Phys. of
Plasmas, **9**, 451 (2002)
- [21] B. Jun, & M.L. Norman, ApJ, **465**, 800 (1996)
- [22] M. Vietri, & L. Stella, ApJL, **507** L45
- [23] A. Königl, & J. Granot, ApJ, **574**, 134 (2002); J. Granot, & D. Guetta, D.
astro-ph/0211136 (2002)
- [24] A.I. MacFadyen & S.E. Woosley, ApJ **524**, 262 (1999)
- [25] S. Akiyama, J.C. Wheeler, D.L. Meier & I. Lichtenstadt, ApJ in press, (2002)
- [26] E.N. Parker, 1993, ApJ, 408, 707 (1993); E.G. Blackman, A. Frank J.A. Markiel,
J.H. Thomas, H.M. & Van Horn, H. M. Nature, **409**, 485 (2001)
- [27] Y. Uchida & K. Shibata PASJ, **37** 31 (1985); J. Contopoulos ApJ **450** 616 (1995);
D. Lynden-Bell , 1996, MNRAS, **279** 389 (1996); S. Heinz & M.C. Begelman ApJ
535 104 (2000); D.A. Uzdensky, ApJ **574** 1011 (2002)
- [28] R.D. Blandford & D.G. Payne MNRAS, 199 883 (1982); R.V.E. Lovelace, J.C.L.
Wang ,M.E. Sulkanen, ApJ, **315** 504 (1987); D.L. Meier, ApJ **522**, 753 (1999)
- [29] N.I. Shakura & R.A. Sunyaev A&A, **24**, 337 (1973)
- [30] S.H. Lubow, J.C.B. Papaloizou, & J.E. Pringle, MNRAS, **267**, 235 (1994)

- [31] S. Kida, S. Yanase, J. & Mizushima, *Physics of Fluids*, **3** 457 (1991)
- [32] J. Maron & S. Cowley, to be submitted to *ApJ* (2002); astro-ph/0111008
- [33] J. Maron & E.G. Blackman, *ApJL* **566**, L41 (2002).
- [34] A. Schekochihin, S. Cowley, J. Maron, J., & L. Malyshkin 2002, *PRE.*, 65, 6305.
- [35] Ya. B. Zeldovich, A.A. Ruzmaikin, D.D Sokoloff, 1983, *Magnetic fields in astrophysics* (Gordon & Breach, New York)
- [36] A. Pouquet, U. Frisch, & J. Leorat, *JFM*, **77**, 321 (1976); N.I. Kleeorin & A.A. Ruzmaikin, *Magnetohydrodynamics*, **2**, 17, (1982)
- [37] E.G. Blackman & A. Brandenburg, in press, *ApJ*, **579**, 359, (2002),
- [38] L. Woltjer, *PNAS*, **44**, 80 (1958).
- [39] M.A. Berger & G.B. Field, *JFM*, **147**, 133 (1984)
- [40] M.A. Berger, *Geophys. & Astrophys. Fluid Dyn.*, **30** 79 (1984).
- [41] N. Seehafer, *PRE*, **53**, 1283 (1996); H. Ji *PRL* **83**, 3198 (1999)
- [42] S.A Balbus & J.F. Hawley, *MNRAS*, **266**, 769 (1996)
- [43] C.A. Tout & J.E. Pringle. *MNRAS* **281** 219 (1996)

Discussion

blackman@pas.rochester.edu Speaker: Eric Blackman

A. Hujeirat, Max Planck Institute: Do you agree that BH-instability is a transient phenomena because the field in sims. never reach $u_A = c_s$? Even if it did, the field will flow up through Parker instability. If you switch on dynamo of Ref. [43] can you produce large scale magnetic fields appropriate for jets?

Blackman: (1) In the saturated (steady, not transient) state, MRI simulations usually show $u_A < c_s$. Since $u_A/c_s \sim \alpha_{ss}^{1/2}$ (see section 4.1) the question of u_A is a question of predicting α_{ss} which is not yet understood, though $\alpha_{ss} \sim 1/2\pi$ is not yet theoretically ruled out (see below, Shibata question). Simulations do however show $u_A \sim u_2$ (turbulent equipartition). (2) The dynamo of [43] might be important (see constraint of sec. 4.1) but in a helical generalization: they discuss the end state of an inverse cascade, not the time evolution. A helical dynamo is needed to make this happen fast enough for GRB.

A. Brandenburg, NORDITA: A crucial question is where the kinetic helicity comes from. You mentioned thermal buoyancy but simulations haven't yet shown this. Instead of expansion of rising tubes we see inflow along field lines toward the field maximum, i.e. contraction and hence the opposite sign of the dynamo α_d .

Blackman: This is indeed important, but fortunately the Poynting flux doesn't care about the sign of α_d as long as its magnitude is not too small. Also, since $\alpha_d \propto \overline{\mathbf{u} \cdot \nabla \times \mathbf{u}} - \overline{\mathbf{b} \cdot \nabla \times \mathbf{b}}$ the current helicity could actually be the main driver in an MRI unstable disk.

K. Shibata, Kyoto Univ.: What is the relation between your theory and the mechanism of MRI field saturation? How do you think this issue will be resolved?

Blackman: As shown in section 4.1, asking how large u_A/c_s gets is also asking how large α_{ss} should be. It is not yet clear why α_{ss} should not be $\sim (1/2\pi)^{1/2}$; one might say that the field produced either by helical or nonhelical dynamos is limited in the thin disk MRI framework only by the strength that keeps the disk thin, or by the strength which for which the vertical Alfvén crossing time equals the orbital period. Since the wavenumber of maximum growth of the MRI satisfies $ku_A \sim \Omega$, as the field energy grows (i.e. as u_A grows), k decreases, but not smaller than that associated with the disc thickness. This would imply $\alpha_{ss}^{1/2} \sim u_A/c_s \sim 1/2\pi$. This argument does not invoke anything about helical large scale dynamo theory except that the ratio could be lowered in practice by large structures as they remove field from a thin disk. Sometimes small boxes show values of $\alpha_{ss} < (1/2\pi)^{1/2}$ but numerical simulations have not converged on a universal value. In thick disks we might expect a higher value than for thin disks, and there are no global simulations of thin disks yet. More theory and simulation are needed to pin this down. It is not even clear that α_{ss} is a constant; it could depend on radius.

S. Kulkarni, Caltech: Focusing on the afterglow external shock, on what time scales can fields grow? I draw your attention to fields in young supernovae (weeks to months old). The inferred B strengths are far in excess of ISM fields. Are these generated by the mechanism(s) you discussed for the external shock or are they compressed fields in the wind of the progenitor?

Blackman: The plasma (two-stream Weibel) instability would saturate on incredibly short time scales, but the associated fields generated are initially on too small a scale without further hydrodynamic evolution. I think the shocks eventually become unstable to MHD turbulence. This turbulence can then further amplify the fields to near equipartition with turbulent motions on a dynamical time scale of the eddies (by the “direct” nonhelical type dynamo discussed) which should be of order a dynamical shock crossing time. Ref. [21] showed that this is promising; fields were amplified to within a factor of 10 of the observed fields. Additional simulations are needed.